

AD-A036 534

CALIFORNIA UNIV SANTA BARBARA DEPT OF PHYSICS  
SPECTROSCOPY IN KINETIC STUDIES OF GAS-PHASE REACTIONS. (U)  
DEC 76 H P BROIDA, R E STEELE

F/G 20/5

AF-AFOSR-2724-74

UNCLASSIFIED

AFOSR-TR-77-0071


NL

| OF |  
AD  
A036534



END

DATE  
FILMED  
3-77



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AFOSR - TR - 77 - 0071

Final Report

~~August 1974 - September 1976~~

1 May

31 Aug.

Organization: AFOSR  
Subelement: Physical ChemistryProject: 9538 Tact: 01  
Grant Number: AFOSR-74-2724

## SPECTROSCOPY IN KINETIC STUDIES OF GAS-PHASE REACTIONS

H. P. Broida  
University of California  
Department of Physics  
Santa Barbara, CaliforniaAbstract

New developments in remote sensing, communications, energy systems and other areas of interest depend upon advances in laser technology. Chemical or transfer laser systems hold promise of fulfilling many of these needs. The development of new and better chemical laser systems, however, awaits increased understanding and knowledge of the reactive and spectral characteristics of candidate molecules. The research under this grant has concentrated on obtaining a detailed understanding of metal-oxidant and metal-halogen reactions. Photon yields and product state branching ratios have been determined for many reactions. In many cases it was necessary to develop basic spectroscopic information on the electronically excited product molecules. In the course of these studies, other molecules of atmospheric, combustion, or chemical laser interest also came under scrutiny, and molecular constants were obtained.

Various diatomic molecular species have been produced in vacuum flow systems; typically metal vapor entrained in an inert gas is reacted with a halogen or oxidant. These exothermic reactions produce product molecules in excited states whose fluorescence (chemiluminescence) is analyzed. Laser induced fluorescence (photoluminescence) is also used to probe both ground and excited electronic states.

The following research summary is organized according to experimental technique.

1472

NO	Attn Section	<input checked="" type="checkbox"/>
CO	Butt Section	<input type="checkbox"/>
TRANSITION		<input type="checkbox"/>
DISCONTINUED		
RE		
DISTRIBUTION/AVAILABILITY NOTES		
DISC	AVAIL. TYPE OR SPECIAL	
A		



462080/CA

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

## 1) Chemiluminescence, Photon Yields, and Branching Ratios

Reactions of lead with  $\text{N}_2\text{O}$ ,  $\text{O}$ ,  $\text{O}_2$ , and  $\text{O}_3$  were investigated under a wide variety of conditions and were found to depend dramatically upon both oxidant species and pressure.<sup>11</sup> Addition of active nitrogen or  $\text{NF}_3$  to the flame resulted in a strong enhancement of the  $\text{PbO}$  band intensity and introduced strong atomic emission while  $\text{CO}$ ,  $\text{NO}$ , and  $\text{N}_2$  quenched the flame. Many new bands were found and molecular constants were determined.

A method was devised to illustrate, on one graph, the population behavior of electronic and vibrational states as the flame conditions were changed. Unless both the electronic and vibrational state populations can be described by a single temperature (and satisfy Boltzmann statistics) the graph will be nonlinear. The technique does not require absolute measurements or knowledge of transition probabilities, but allows use of direct relative photomultiplier signals. The various oxidants used in the  $\text{PbO}$  experiment gave widely varying population distributions in the product molecule.

Modifications to the standard furnace system allowed extension of its upper temperature limit by several hundred degrees, to 2100 K, and permitted production of  $\text{TiO}$ .<sup>14</sup>  $\text{O}_2$ ,  $\text{N}_2\text{O}$ ,  $\text{NO}_2$ ,  $\text{NO}$ , and  $\text{CO}_2$  were used as oxidants of the titanium atoms; the first two species were chosen for detailed study.  $\text{Ti} + \text{O}_2$  flames had electronic and vibrational temperatures of  $2500 \pm 500$  K, while  $\text{Ti} + \text{N}_2\text{O}$  flames were approximately twice as hot, and exhibited wider departures from Boltzmann statistics. Ultraviolet emission was three orders of magnitude higher with  $\text{N}_2\text{O}$  than with  $\text{O}_2$ .

Confirmation of our matrix isolation determination<sup>4</sup> of the  $a^1\Delta - X^3\Delta$  spacing as  $3500 \text{ cm}^{-1}$  was obtained from intensity data. Molecular constants of the  $E^3\Pi$  state were determined. An anomalous feature in the  $\text{TiO}$  spectrum was attributed to  $\text{YO}$  contamination and preliminary work on a study of yttrium reacting with oxygen has commenced.

Preliminary photon yield data have been taken for the reactions of copper with  $\text{F}_2$ ,  $\text{NF}_3$  and  $\text{SF}_6$ . Chemiluminescence spectra of  $\text{CuF}$  have been taken in the 200-1100 nm region and corrected by computer for relative spectral response of the detection system. Three new electronic states have been discovered with transitions at 438, 568 and 680 nm; strong perturbations in the spectrum of the  $A^1\Pi - X^1\Sigma$  transition are caused by the close proximity of the upper state in the new 568 nm transition. Total photon yields from 200-800 nm were less than 0.5% for the reaction  $\text{Cu} + \text{F}_2$ , and lower for  $\text{Cu} + \text{NF}_3$ ; 60-90% of the emission was in the A-X transition, and 10-40% divided between the B-X and C-X systems.

Aluminum atoms diluted in a buffer gas were reacted with  $\text{F}_2$ ,  $\text{Cl}_2$ ,  $\text{Br}_2$ ,  $\text{I}_2$ ,  $\text{NF}_3$ , and  $\text{SF}_6$  to produce diatomic aluminum halides in chemiluminescent flames.<sup>13</sup> Emission from  $a^3\Pi$  to  $X^1\Sigma^+$  was observed for all the halides, and for  $\text{Al}$ ,  $\text{AlF}$  and  $\text{AlCl}$ , higher electronic states were also observed. Photon yields were 2% for  $\text{Al} + \text{NF}_3$ , and less than 0.01% for the other reactants. In the case of  $\text{Al} + \text{NF}_3$ , about 90% of the total yield was due to  $\text{Al}$  atomic emission and in fact the atomic emission extended up to the ionization limit at 207 nm. Energy transfer from metastable molecules was invoked as the excitation mechanism.

\*References and talks, listed in chronological order, are found following section 9.



Observation of intercombination emission from AlF allowed the first direct determination of the  $a^3\Pi - X^1\Sigma$  separation,<sup>9</sup> which agreed well with an earlier value calculated from vibrational perturbations observed in the  $A^1\Pi$  state.

Emission from ultraviolet to near infrared was observed from flames of Al atoms reacting with various oxidants.<sup>3</sup> The species  $N_2O$ ,  $O_2$ , microwave discharged  $O_2$ ,  $O_3$ ,  $CO_2$ ,  $NO_2$ ,  $NO$  and  $CO$  were used to form the  $AlO$ . Photon yields for the reactions varied widely from 3% for  $Al +$  microwave discharged  $O_2$ , and 2% for  $O_3$ , to 0.4% for  $N_2O$  and less than 0.005% for all other oxidants. The  $A^2\Pi$  to  $X^2\Sigma$  system, which had not previously been observed in emission, was seen with several of the oxidants. A broad band emission feature present with excess oxidant was attributed to further oxidation products of  $AlO$ .

An isotopic substitution experiment<sup>10</sup> of Ba reacting with  $N_2^{16}O$  or  $N_2^{18}O$  allowed unique absolute numbering of the vibrational states in  $BaO$ , and subsequent determination of the spectroscopic constants of the  $A^1\Pi$  state.

## 2) Photoluminescence

A single mode tunable dye laser was used to induce photoluminescence from  $CuF$ . High resolution scans near the origin of the  $A^1\Pi - X^1\Sigma$  transition show strong perturbations from a newly discovered state at around  $17600\text{ cm}^{-1}$ . This perturbation is exhibited as an extremely large  $\Lambda$ -type doubling of the  $A^1\Pi$  state ( $q \approx .01\text{ cm}^{-1}$ ).

Six wavelengths of the  $Ar^+$  laser have been found to be coincident with transitions in  $Cu_2$ . Five of these lines ranging from  $457.9\text{ nm}$  to  $488.0\text{ nm}$  excite the  $B^1\Sigma_u$  state, while the  $496.5$  line excites the  $A^1\Pi$  state.

Laser induced photoluminescence has been observed from  $TiO$  molecules<sup>15</sup> produced in a chemiluminescent flame of  $Ti + O_2$ .  $Ar^+$  and cw dye lasers were used in a detailed study of the  $\alpha(C^3\Delta - X^3\Delta)$  system in emission. Techniques were developed for assigning photoluminescence transitions involving states of high multiplicity in molecules with several isotopes. Emission in the  $0-0$  band of the  $\beta(c^1\phi - a^1\Delta)$  system, observed with  $476.5\text{ nm}$  excitation, was shown to result from collisional relaxation of the  $C^3\Delta(v=2)$  to  $c^1\phi(v=0)$  states. Assignment of a new  $C^3\Delta_3 - a^1\Delta$  transition places the  $a^1\Delta$  state at  $3444 \pm 10\text{ cm}^{-1}$  above the  $X^3\Delta$  state, thus confirming and refining earlier measurements made in this laboratory of the singlet-triplet separation. These experiments have greatly improved the overall spectroscopic understanding of this astrophysically important molecule.

## 3) Lifetime Measurements

The radiative lifetimes have been measured for the  $A^1\Pi$ ,  $B^1\Sigma$  and  $C^1\Pi$  states of  $CuF$  and have been found to be .5, .7 $\mu$ s, and 12 microseconds respectively. Tunable pulsed dye lasers were used for the measurements with either flashlamp (1  $\mu$ sec duration) or nitrogen laser (1 nsec pulse duration) excitation. A computerized detection system was developed to enable signal averaged and wavelength scanned lifetime measurements to be made. A high resolution time-resolved spectroscopic study of the  $A^1\Pi$  state presently in progress is expected to elucidate the detailed nature of this highly perturbed state.

Radiative lifetimes have been measured<sup>12</sup> for the A and B states of  $Cu_2$ . Using a  $N_2$ -pumped tunable dye laser, lifetimes of  $30 \pm 15\text{ nsec}$  at a background argon pressure of .5 torr and  $20 \pm 15\text{ nsec}$  at 10 torr were measured for the first three vibrational levels of the B state. The A state lifetime of the  $v=0$  level is  $70 \pm 15\text{ nsec}$  at .5 torr and  $50 \pm 15\text{ nsec}$  at 10 torr.

#### 4) Computer Analysis

During the period of this grant a dedicated computer data handling system was purchased through another funding source. This system has proved itself a great utility in correcting spectra for instrument response, and in comparing calculated to observed spectra.

In studies of the  $\text{CuF}$ ,  $A^1\Pi$  state, comparison of data to synthetic spectra revealed the necessity of using different B values for the Q vs the P and R branches to obtain suitable agreement, and thereby revealed the existence of a new state which was responsible through perturbations for the anomalies in the  $A^1\Pi$  state.

Such comparison of synthetic and observed spectra also facilitates determination of rotational and vibrational temperatures in chemiluminescent flames.

#### 5) Excitation Transfer

Combustion of Mg vapor in  $\text{N}_2\text{O}$  with He and  $\text{CO}_2$  diluents in a low-loss laser cavity resulted in cw oscillation at  $10.6\text{ }\mu\text{m}$ .<sup>7,5</sup> The pumping takes place via chemical reaction and subsequent electronic-vibrational (E-V) energy transfer. The first step, whose exothermicity ultimately drives the laser, is the formation of electronically excited  $\text{MgO}^*$ . A large proportion of this  $\text{MgO}^*$  is in states whose radiative decay is slow. In the presence of  $\text{CO}_2$  the dominant mode of destruction of these metastable  $\text{MgO}^*$  molecules is believed to be collisional conversion of their electronic excitation to vibrational excitation of  $\text{CO}_2$ , with the consequent inversion of the (001) level with respect to the (100) level.

Further studies of this topic under support of AFOSR grant 76-2959 demonstrated that the population of  $\text{MgO}^*$  will react efficiently with CO to yield  $\text{Mg}(^3\text{P})$  atoms and excited  $\text{CO}_2$ .

High-lying Rydberg levels of Al were excited<sup>8</sup> in a chemical reaction with  $\text{NF}_3$ . A possible mechanism is that high lying levels of Al are populated by energy transfer from excited molecular nitrogen produced in the  $\text{Al} + \text{NF}_3$  mixture. Al atomic states up to the series limit were observed with  $\text{NF}_3$  as well as in another experiment with microwave discharged  $\text{N}_2$ . While with active nitrogen, the resulting populations of the Al excited states followed a Boltzmann distribution at about 3000 K, in the case of the  $\text{NF}_3$  the population distribution was non-Boltzmann and skewed toward higher energies.

Active nitrogen was also used in experiments on  $\text{TiO}$  and  $\text{PbO}$  flames,<sup>11,14</sup> where it materially enhanced the molecular emission as well as causing intense atomic emission from the metal. In the case of Pb the effect was primarily to increase the electronic temperature while maintaining a Boltzmann distribution. It is believed that  $\text{PbO}^*$  was produced by transfer from active nitrogen to  $\text{PbO}$ . Similar explanations apply in the case of  $\text{TiO}$ .

#### 6) Gain Measurements

Intracavity laser gain measurements using a  $\text{CO}_2$  electrical discharge probe laser were made to study flames of Ba, Ca and Mg burning in a mixture of  $\text{N}_2\text{O}$ ,  $\text{CO}_2$  and He.<sup>5</sup> Gain at  $10.6\text{ }\mu\text{m}$  was detected for each of these metals. For the Mg flames the gain coefficient was found to be in the range of  $10^{-3} - 10^{-2}/\text{cm}$ . The addition of  $\text{N}_2$  diluent enhanced the gain, whereas CO diluent quenched it.

These measurements led to the observation of cw lasing as mentioned above.

# Publications

1. J. M. Brom, Jr.  
H. P. Broida      Photodissociative Production of  $O(^1S)$  and  $N(^2D)$  From  $N_2O$  in an Argon Matrix at 4 K, Chem. Phys. Lett. 33, 384 (1975).
2. M. M. Hessel  
R. E. Drullinger  
H. P. Broida      Chemiluminescent Reactions in a Heat Pipe Oven  
J. Appl. Phys. 46, 2317 (1975).
3. S. Rosenwaks  
R. E. Steele  
H. P. Broida      Chemiluminescence of  $AlO$ , J. Chem. Phys. 63, 1963 (1975).
4. J. M. Brom, Jr.  
H. P. Broida      Laser Photoluminescence of  $TiO$  in Ne at 4 K, J. Chem. Phys. 63, 3718 (1975).
5. D. J. Benard      CW Chemical Laser Gain at 10.6 Microns, Chem. Phys. Lett. 35, 167 (1975).
6. R. W. Field  
G. A. Capelle  
M. A. Revelli      Optical-Optical Double Resonance Laser Spectroscopy of  $BaO$   
J. Chem. Phys. 63, 3228 (1975).
7. D. J. Benard      CW Chemical Transfer  $CO_2$  Laser, Appl. Phys. Lett. 27, 542 (1975).
8. S. Rosenwaks  
H. P. Broida      Chemiexcitation Transfer to High Lying Rydberg Levels of  $Al$   
J. Opt. Soc. Am. 66, 75 (1976).
9. S. Rosenwaks  
R. E. Steele  
H. P. Broida      Observation of  $a^3\Pi - X^1\Sigma$  Intercombination Emission in  $AlF$   
Chem. Phys. Lett. 38, 121 (1976).
10. J. C. Wyss  
H. P. Broida      Vibrational Analysis of the  $A'^1\Pi$  State of Barium Oxide Using Two Isotopes, J. Mol. Spectros. 59, 235 (1976).
11. C. Linton  
H. P. Broida      Chemiluminescent Spectra of  $PbO$  from Reactions of  $Pb$  Atoms  
J. Mol. Spectros. 62, 396 (1976).
12. R. E. Steele      Photoluminescence, Lifetimes, and Discharge Excitation of  $Cu_2$   
J. Mol. Spectros. 61, 477 (1976).
13. S. Rosenwaks      Chemiluminescent Reactions of  $Al$  Atoms and Halogens  
J. Chem. Phys. 3668 (1976).
14. C. Linton  
H. P. Broida      Flame Spectroscopy of  $TiO$ : I. Chemiluminescence  
J. Mol. Spectros. xx, (in press 1976).
15. C. Linton  
H. P. Broida      Flame Spectroscopy of  $TiO$ : II. Photoluminescence of the  $\alpha(C^3\Delta - X^3\Delta)$  System, J. Mol. Spectros. xx, (in press 1976).



Talks and Presented Papers

1. 10-74 4th Conference on Chemical and Molecular Lasers,  
St. Louis, Missouri  
"Chemiluminescence and Photoluminescence Spectroscopy of  
the Barium Monohalides" (R. S. Bradford, Jr.)
2. 10-74 4th Conference on Chemical and Molecular Lasers,  
St. Louis, Missouri  
"The Diatomic Oxides and Halides of Ca and Sr: Photon  
Yields and  $A^1\Pi$  Molecular Constants" (G.A. Capelle)
3. 10-74 Physics Department Colloquium, University of California,  
Santa Barbara, CA.  
"A Recent Visit to Laser Science in the Soviet Union"  
(H. P. Broida)
4. 10-74 Seminar, Stanford Research Institute, Menlo Park, CA.  
"A Recent Visit to Laser Science in the Soviet Union"  
(H. P. Broida)
5. 11-74 Seminar, NBS Washington, D. C.  
"A Recent Visit to Laser Science in the Soviet Union"  
(H. P. Broida)
6. 12-74 Physics Seminar, University of California, Santa Barbara  
"An Attempt to Measure Atmospheric NO Density"  
(H. P. Broida)
7. 1-75 American Physical Society Meeting, Anaheim, CA  
"Red ( $A^2\Pi-X^2\Sigma$ ) System of the AlO Molecule",  
(R. E. Steele)
8. 2-75 University of Fredericton, New Brunswick Canada,  
"Chemiluminescence and Photoluminescence of Diatomic Molecules"  
(H. P. Broida)
9. 3-75 University Federal de Pernambuco, Department of Physics, Brazil  
"Chemiluminescence and Photoluminescence of Diatomic Molecules"  
(H. P. Broida)
10. 3-75 University Federal de Pernambuco, Brazil  
"Flame Spectroscopy as a Research Tool"  
(H. P. Broida)
11. 3-75 University of Denver, Chemistry Department, Colorado  
"Flames"  
(H. P. Broida)
12. 9-75 Second Summer Colloquium on Electronic Transition Lasers,  
Woodshole Mass.  
"Sensing Chemically Excited Metastable Populations by CO<sub>2</sub>  
Laser Gain Measurements"  
(D. J. Benard)

13. 9-75 Second Summer Colloquium on Electronic Transition Lasers, Woodshole, Mass.  
"Chemiexcitation Transfer from Metastable Molecules to Metal Atoms in Metal-Halogen Flames"  
(S. Rosenwaks)
14. 10-75 Molecular Spectroscopy Meeting, UCSB, Santa Barbara, CA,  
"A Summer of Frustration"  
(H. P. Broida)
15. 11-75 Annual Meeting of the Division of Atomic and Molecular Physics, Canadian Association of Physicists, Fredericton, New Brunswick, Canada  
"Optical Spectroscopy of Metal Atom Flames"  
(H. P. Broida)
16. 12-76 American Physical Society Meeting, Stanford, CA  
"Chemiluminescence and Photoluminescence of Gas Phase CuF"  
(R. E. Steele, to be given).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>18</b> AFOSR - <b>19</b> TR - 77 - 0071	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>6</b> SPECTROSCOPY IN KINETIC STUDIES OF GAS-PHASE REACTIONS.	5. TYPE OF REPORT & PERIOD COVERED <b>9</b> Final <i>rept.</i> 1 May 74 - 31 Aug 76	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) <b>10</b> H. P. Broida, R. E. Steele	8. CONTRACT OR GRANT NUMBER(s) AFOSR-74-2724	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics University of California Santa Barbara, CA 93106 <b>11</b> Dec 76	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 9538-01 61102F 681303	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (NC) Bolling AFB, Bldg, 410 Bolling, AFB, D.C. 20323	12. REPORT DATE December 1976	13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>12</b> 9p.	15. SECURITY CLASS. (of this report) UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
<b>15</b> ✓ AF-AFOSR-2724-74		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <b>16</b> 9538 / <b>17</b> 01		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) New developments in remote sensing, communications, energy systems and other areas of interest depend upon advances in laser technology. Chemical or transfer laser systems hold promise of fulfilling many of these needs. The development of new and better chemical laser systems, however, awaits increased understanding and knowledge of the reactive and spectral characteristics of candidate molecules. The research		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

400628

B

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

under this grant has concentrated on obtaining a detailed understanding of metal-oxidant and metal-halogen reactions. Photon yields and product state branching ratios have been determined for many reactions. In many cases it was necessary to develop basic spectroscopic information on the electronically excited product molecules. In the course of these studies, other molecules of atmospheric, combustion, or chemical laser interest also came under scrutiny, and molecular constants were obtained.

Various diatomic molecular species have been produced in vacuum flow systems; typically metal vapor entrained in an inert gas is reacted with a halogen or oxidant. These exothermic reactions produce product molecules in excited states whose fluorescence (chemiluminescence) is analyzed. Laser induced fluorescence (photoluminescence) is also used to probe both ground and excited electronic states.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



